AGS STUDIES REPORT

NUMBER 94

Date 8/8,9,11,15,26,29,31/77

Experimenters E. Raka

Subject Vertical Beam Size and Emittance at Low and High Intensity

OBSERVATIONS AND CONCLUSION

<u>Purpose</u>: To measure the vertical emittance of the AGS beam at different intensity levels and momenta in order to determine the magnitude of phase space dilutions that are present.

AGS Conditions: Measurements were made parasitically on SEB runs at 5 GeV and $\approx 3 \times 10^{12} \ \rm protons/pulse$ and normal 28.5 GeV SEB running at $\approx 9.5 \times 10^{12}/\rm pulse$; linac current $\approx 70 \ \rm mA$. $\approx 110 \ \rm \mu sec$ at high intensity and 5-6 turns at low intensity.

<u>Procedure</u>: The J19 vertical targets were used to remove either 5 of 10% of the beam from the top or bottom. Records of the beam frequency and gauss clock reading were made at the end of the loss pattern as observed on the normalized current transformer. The latter signal was measured before and after by an A to D converter and the difference multiplied by five displayed.

Results:

	Size	βγ	GC	Emittance*	Time	
3x10 ¹²	1.45"	1.33	~-	20.5 πμradM	95 msec	
6 turns 79 mA	.77''	5.34		23.2	175 ''	
3X10 ¹² 5 turns 76 mA	1.215"	1.34		14.5* "	96 ''	*90% emittance
9.5×10 ¹²	1.675"	1.39	2459	28.5 "	94 ''	
72 mA	.845"	6.88	12625	36 "	194 ''	
	.795"	6.88	11	31.8 "*	194* "	*zero theta skew
1.3×10 ^{12*}	1.88"*	.686		17.7 "°	52*	quad program on. *i.e. injection spiraling beam only

95% except as noted.

Comments: Both the low and high intensity results indicate that the emittance growth from about twice injection momentum to near transition energy can be controlled. There is still significant growth from injection to a $\beta\gamma\approx 1.4$ although the accuracy of the spiraling beam measurement is less than that of the others. Also HEBT emittance of $\approx 5~\pi$ for 90% have been obtained by J-L LeMaire which imply considerable dilution in the injection process. The latter is apparently only mildly intensity dependent.